



Contents lists available at ScienceDirect

Clinical Nutrition

journal homepage: <http://www.elsevier.com/locate/clnu>

Opinion paper

Modulation of Hb-O₂ affinity to improve hypoxemia in COVID-19 patientsSimon Woyke^a, Simon Rauch^{b,c}, Mathias Ströhle^d, Hannes Gatterer^{b,*}^a Department of Anesthesiology and Intensive Care Medicine, Medical University of Innsbruck, Innsbruck, Austria^b Institute of Mountain Emergency Medicine, Eurac Research, Bolzano, Italy^c Department of Anesthesia and Intensive Care Medicine, "F. Tappeiner" Hospital, Merano, Italy^d Department of General and Surgical Critical Care Medicine, Medical University of Innsbruck, Innsbruck, Austria

ARTICLE INFO

Article history:

Received 15 April 2020

Accepted 24 April 2020

Keywords:

COVID-19

Nutritional supplement

5-Hydroxymethylfurfural

SUMMARY

This opinion paper aims at discussing the potential impact of modulating the Hb-O₂ affinity by the nutritional supplement 5-HMF on patients affected by COVID-19. The paper describes the critical role of the oxygen affinity in hypoxemic COVID-19 patients and the potential positive effect of 5-HMF, a compound shown to increase the Hb-O₂ affinity.

© 2020 Elsevier Ltd and European Society for Clinical Nutrition and Metabolism. All rights reserved.

1. Introduction and discussion

The oxygen dissociation curve (ODC) describes the dependency of the oxygen saturation on the oxygen partial pressure (PO₂) [1]. With its sigmoid shape, the curve is subjected to right or left shifts, thereby changing hemoglobin-O₂ affinity. An increase in the partial pressure of carbon dioxide (PCO₂), in 2,3-diphosphoglycerate (2,3 DPG) and in temperature as well as a decrease in pH lead to a right-shift of the ODC, decreasing oxygen affinity, and vice versa [1,2]. In hypoxemic conditions, a shift of the ODC can significantly alter arterial oxygen saturation (SaO₂) and consequently arterial oxygen content (CaO₂). With a PO₂ of 60 mmHg, for instance, SaO₂ may be around 81% [1] at a pH of 7.27 (corresponding to a PCO₂ of 60 mmHg according to Siggaard-Andersen nomogram) and normal body temperature (37 °C), or about 90% at a pH of 7.4, 37 °C body temperature and a PCO₂ of 40 mmHg [3]. Assuming a hemoglobin concentration of 12 g/dl, this corresponds to an increase in CaO₂ from 13.0 to 14.5 ml/dl. Further assuming a cardiac output of 6 l/min, the delivery of oxygen (DO₂) increases from 780 to 870 ml/min. This increase in DO₂ by augmenting CaO₂ is equivalent to a DO₂ increase by rising cardiac output by approximately 0.7 l/min. This approximates the effect of low- to medium-dose Dobutamine, a commonly used inotrope [4]. The difference in SpO₂ may even widen with lower PO₂, due to the shape of the ODC, or

changes in body temperature, leading to an even greater effect on DO₂ as outlined before.

Coronavirus disease 2019 (COVID-19) is a respiratory tract infection caused by a newly emergent coronavirus [5]. The disease is characterized by symptoms of cough and high fever and causes primarily interstitial lung changes and is characterized by a dissociation between relatively well preserved lung mechanics and the severity of hypoxemia [5]. Intrapulmonary ventilation-perfusion mismatch, probably due to impaired hypoxic vasoconstriction, results in hypoxemic respiratory failure [6]. Oxygen administration via face mask or nasal cannula, high-flow-nasal oxygen or non-invasive ventilation are common first-line interventions to improve oxygenation and dyspnea in these patients [5]. In case of persistent hypoxemia or clinical deterioration, endotracheal intubation and invasive ventilation is required. In mechanically ventilated adults with COVID-19, gentle ventilation with low tidal volume is recommended in order not to further damage the lungs [7,8]. Low tidal volume ventilation often results in hypercapnia, which is tolerated to pH levels as low as 7.2 (permissive hypercapnia) [9]. As outlined above, the consequence of the high PCO₂ and increased body temperature is a shift of the ODC to the right, worsening hypoxemia. In these patients, the PO₂ levels are on the steep part of the ODC, so that a small shift of the ODC may have a large effect on the oxygen affinity and thus on SpO₂. The question arises whether shifting the ODC back to the left would improve oxygenation of COVID-19 patients.

5-hydroxymethylfurfural (5-HMF) is an agent able to shift the oxygen dissociation curve to the left. 5-HMF reduces the P50 (PO₂

* Corresponding author.

E-mail address: hannes.gatterer@eurac.edu (H. Gatterer).

at which 50% of hemoglobin is saturated with oxygen, a parameter indicating the position of the ODC) via allosteric modification of the hemoglobin [10]. Animal studies showed that under severe hypoxia conditions (FiO_2 : 5%), 5-HMF increased hemoglobin affinity, preserved systemic O_2 delivery and partially was able to maintain microvascular oxygenation (i.e., by protecting arteriolar and venular vasodilation and blood flow) [10]. In swine exposed to severe hypoxia, 5-HMF treatment decreased P50, improved SaO_2 , and mitigated increases in pulmonary artery pressure [11]. Moreover, a recent study indicated that the substance might have cardiac protective properties by inhibiting L-type Ca^{2+} channels [12]. However, not only in animal and in vitro models beneficial effects were reported. 5-HMF was also found to increase the oxygen affinity in healthy subjects exposed to hypoxia and in sickle cell disease patients [13,14]. Additionally, the combined oral intake of 5-HMF and α -ketoglutaric acid increased SpO_2 during cycling exercise at 3500 m in healthy subjects [15]. The oral intake of these 2 substances was also shown to reduce oxidative stress, to increase exercise capacity and to reduce ICU and hospitalization time in patients admitted for lung resection [16]. It is important to mention that these substances are available as nutritional supplements and that no relevance for humans concerning carcinogenic and genotoxic effects have been found for 5-HMF supplementation [17].

This makes 5-HMF a potential therapeutic agent for the treatment of hypoxemic COVID-19 patients. By facilitating oxygen loading and increasing SaO_2 , breathing exertion might be reduced. This could delay or even prevent invasive ventilation and hence save valuable intensive care capacity in a crisis like this. In ARDS patients undergoing lung protective ventilation with permissive hypercapnia, a left-shift of the ODC induced by 5-HMF might ameliorate oxygenation by counteracting the hypercapnia effect on the oxygen affinity. On the other hand, a left-shift of the ODC might impair peripheral O_2 unloading, leading to an unchanged cellular oxygen supply despite an increased DO_2 . Even though comparability might be questioned, recent data on hypoxic exercise show that a high O_2 -affinity and hence enhanced oxygen uptake in the lungs, outweighed deficits in peripheral O_2 unloading [2,18]. Additionally, it was reported previously that in hypoxic conditions O_2 unloading from Hb does not require a right-shift of the ODC [19].

In conclusion, the modulation of hemoglobin-oxygen affinity by 5-HMF might be worth investigating as a potential therapeutic target in hypoxemic respiratory failure, e.g. due to COVID-19.

Conflict of interest

None.

References

- [1] Mairbaurl H, Weber RE. Oxygen transport by hemoglobin. *Comp Physiol* 2012;2:1463–89.
- [2] Dempsey JA. With haemoglobin as with politics - should we shift right or left? *J Physiol* 2020;598:1419–20.
- [3] Mairbaurl H. Red blood cell function in hypoxia at altitude and exercise. *Int J Sports Med* 1994;15:51–63.
- [4] Egstrup M, Gustafsson I, Andersen MJ, Kistorp CN, Schou M, Tuxen CD, et al. Haemodynamic response during low-dose dobutamine infusion in patients with chronic systolic heart failure: comparison of echocardiographic and invasive measurements. *Eur Heart J Cardiovasc Imag* 2012;14:659–67.
- [5] Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395:497–506.
- [6] Gattinoni L, Coppola S, Cressoni M, Busana M, Chiumello D. Covid-19 does not lead to a "typical" acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2020. <https://doi.org/10.1164/rccm.202003-0817LE>.
- [7] Poston JT, Patel BK, Davis AM. Management of critically ill adults with COVID-19. *JAMA* 2020. <https://doi.org/10.1001/jama.2020.4914>.
- [8] Alhazzani W, Moller MH, Arabi YM, Loeb M, Gong MN, Fan E, et al. Surviving sepsis campaign: guidelines on the management of critically ill adults with coronavirus disease 2019 (COVID-19). *Crit Care Med* 2020. <https://doi.org/10.1097/CCM.0000000000004363>.
- [9] Griffiths MJD, McAuley DF, Perkins GD, Barrett N, Blackwood B, Boyle A, et al. Guidelines on the management of acute respiratory distress syndrome. *BMJ Open Respir Res* 2019;6:e000420.
- [10] Yalcin O, Cabrales P. Increased hemoglobin O_2 affinity protects during acute hypoxia. *Am J Physiol Heart Circ Physiol* 2012;303:H271–81.
- [11] Mahon RT, Ciarlone GE, Roney NG, Swift JM. Cardiovascular parameters in a swine model of normobaric hypoxia treated with 5-Hydroxymethyl-2-Furfural (5-HMF). *Front Physiol* 2019;10:395.
- [12] Wolkart G, Schrammel A, Koyani CN, Scherubel S, Zorn-Pauly K, Malle E, et al. Cardioprotective effects of 5-hydroxymethylfurfural mediated by inhibition of L-type Ca^{2+} currents. *Br J Pharmacol* 2017;174:3640–53.
- [13] Stern W, Matthews D, McKew JC, Shen J, GJ K. A phase 1, first-in-man, dose–response study of Aes-103 (5-HMF), an anti-sickling, allosteric modifier of hemoglobin oxygen affinity in healthy normal volunteers. *Blood* 2012;120:3210.
- [14] Hannemann A, Cytlik UM, Rees DC, Tewari S, Gibson JS. Effects of 5-hydroxymethyl-2-furfural on the volume and membrane permeability of red blood cells from patients with sickle cell disease. *J Physiol* 2014;592:4039–49.
- [15] Kossler F, Mair L, Burtscher M, Gatterer H. 5-Hydroxymethylfurfural and alpha-ketoglutaric acid supplementation increases oxygen saturation during prolonged exercise in normobaric hypoxia. *Int J Vitam Nutr Res* 2019;1–6.
- [16] Matzi V, Lindenmann J, Muench A, Greilberger J, Juan H, Wintersteiger R, et al. The impact of preoperative micronutrient supplementation in lung surgery. A prospective randomized trial of oral supplementation of combined alpha-ketoglutaric acid and 5-hydroxymethylfurfural. *Eur J Cardio Thorac Surg* 2007;32:776–82.
- [17] Abraham K, Gurtler R, Berg K, Heinemeyer G, Lampen A, Appel KE. Toxicology and risk assessment of 5-Hydroxymethylfurfural in food. *Mol Nutr Food Res* 2011;55:667–78.
- [18] Dominelli PB, Wiggins CC, Baker SE, Shepherd JRA, Roberts SK, Roy TK, et al. Influence of high affinity haemoglobin on the response to normoxic and hypoxic exercise. *J Physiol* 2020;598:1475–90.
- [19] Calbet JA, Losa-Reyna J, Torres-Peralta R, Rasmussen P, Ponce-Gonzalez JG, Sheel AW, et al. Limitations to oxygen transport and utilization during sprint exercise in humans: evidence for a functional reserve in muscle O_2 diffusing capacity. *J Physiol* 2015;593:4649–64.